



## Getting the Most out of Your Vegetable Garden Soil Test Report

1. Be certain your soil pH is in the optimal range.
2. Add a nitrogen (N) source at the recommended rate.
3. Add a phosphorus (P) or potassium (K) source only if soil test levels are low.
4. Organic matter will reduce the amount of nitrogen needed.

### Soil pH in the 6.0 to 7.0 range is optimal for the health of most vegetable plants.

When soil pH is outside this optimal range, lime can be mixed in to raise soil pH. Sulfur or acidifying fertilizers (like ammonium sulfate or ammonium nitrate) can be mixed in to lower soil pH.

The recommended rate of material should be mixed into the upper 4 to 6 inches of soil. To avoid harming plant roots and to provide the necessary time for the material to adjust the pH throughout the root zone, it is best to mix liming or acidifying material into the soil in the fall after harvest.

When a large amount of material is recommended, such as 20 pounds per 100 square feet, mix half in the fall and the other half in the spring before planting.

### Promote plant health, increase yield, and save money with proper fertilizer applications.

Nutrients required by vegetables in the highest quantity are nitrogen (N), phosphorus (P) and potassium (K). Other essential nutrients are usually available in adequate quantities when the pH is in the optimal range. There is no benefit to applying more fertilizer than plants require. Over application of nutrients may harm plant growth and the environment.

### Fertilizer Sources

Organic matter is added to the soil to improve soil properties by increasing the water and nutrient holding capacity, improving aeration and drainage, feeding microorganisms and providing some nutrients. Common organic matter sources include composted manure, peat moss, plant-based compost and plant and animal by-products.

**Animal manures and plant-based compost** (yard waste, grass clippings, food waste) are readily available and popular fertilizers commonly considered as *natural organic*.

Nutrient content of composted products will vary depending on the source (plant or animal), moisture content, how it was stored and how long it was composted. Weed seeds, a high salt content and pathogens might also be a problem with fresh manure or other organic materials if they are not fully composted. Knowing the nutrient and salt content of your compost or manure pile would require testing every time one of the above variables changed so it is not practical.

The nutrient analysis of commercially available manure and compost is found on the bag. Keep in mind these products may also have a high salt content and only 5 to 20% of the nitrogen will be available to the plants during the first year of application.

Mix in only manure or other materials that are fully composted. Fully composted manure has aged for at least 6 to 9 months. Incorporate into the upper 4 to 6 inches of soil after harvest and before planting to allow time for the slow-release of nutrients in the root zone and to minimize runoff.

**Too much of a good thing?** *If a regular soil test indicate nutrient levels exceed crop needs, limit compost applications to promote healthy soil for plant success.*

**Annually replenishing organic matter to improve soil health is a common practice.** Sandy soils with an organic matter content of 2-2½% and clayey soils with 3-5% organic matter will perform fine. **Organic matter levels twice this amount could lead to nutrient imbalances.**

Keep in mind some nitrogen will be released as soil microorganisms break down organic matter in soil. For every 1% organic matter on the soil test report, you can reduce your applied nitrogen 0.5 ounce per 100 square feet for the year.

**Inorganic** fertilizers are compounds that are chemically synthesized from basic raw materials, isolated from naturally occurring sources or mined materials that provide nutrients. They are less expensive and have a higher nutrient content. They are water-soluble and are quickly available to plants, easy to apply and especially useful in cool weather.

Care should be taken when using *water-soluble fertilizer* sources because they are made of salts that can burn plants. If over applied and watered in excessively they can leach beyond the root zone and end up in the ground water which impacts water quality negatively.

**The nutrient content (fertilizer analysis) is required on every bag of commercially available fertilizer.** The first number indicates the *percent* of nitrogen (N), the second number is the percent of phosphate (P<sub>2</sub>O<sub>5</sub>) a source of phosphorus, and the third number is the percent of potash (K<sub>2</sub>O) a source of potassium. They are simply referred to as N-P-K. A 50 lb. bag of 5-10-5 contains 2½ lbs. of N, 5 lbs. of P<sub>2</sub>O<sub>5</sub> and 2½ lbs. of K<sub>2</sub>O. The rest of the material is made up of other inert material, such as sand or clay granules to help spread the fertilizer.

When nitrogen (N), phosphorus (P) and potassium (K) are all needed, use a *complete fertilizer* that contains all 3 nutrients. For example: 10-10-10, 5-10-5, or 5-10-10. Use Table E to determine which complete fertilizer could be used based on your soil test results.

**Recommendations for phosphorus (P) or potassium (K) nutrients will only be made if nutrients are low.** Additional P or K should not be added unless soil test levels are low.

### **Timing of fertilizer applications**

*Pre-plant* fertilizing involves mixing fertilizer into the upper 4 to 6 inches of the soil before plants are in the ground. It can effectively address soil deficiencies or replenish nutrients absorbed by plants root systems in previous seasons.

*Starter solutions* are liquid fertilizers generally containing a small amount of nitrogen (N) and potassium (K) and a high amount of phosphorus (P) such as 4-12-4 or 10-52-10 and are commonly used when setting out vegetable transplants. Half the rate suggested on the commercial label is recommended.

*Side dressing* refers to the addition of fertilizer during the growing season. The timing varies by crop. See details in Table B.

## Nitrogen (N)

The nitrogen content of a soil sample does not necessarily reflect the future availability of nitrogen because unlike phosphorus and potassium, the nitrogen level will fluctuate depending on biological activity and soil conditions. Nitrogen is one of the most important nutrients for plant growth and is needed in relatively large amounts by all plants so a standard nitrogen recommendation is routinely provided for vegetable gardens. The annual recommended nitrogen rate is 3.5 ounces per 100 square feet or 2.2 pounds per 1,000 square feet.

Organic matter such as aged animal manures and plant-based compost can supply some nitrogen throughout the growing season, however, nitrogen may not be released from these sources early in the season. When the soil temperature is below 50 degrees Fahrenheit, the microorganism activity that breaks down organic matter to release nitrogen is minimal.

The organic matter in your soil is listed on your soil test report. For every 1% organic matter in your soil, you can reduce the recommended nitrogen rate by 0.5 ounce per 100 square feet for the year. Most gardens have at least 2% organic matter so you will only need to apply 2.5 ounces of nitrogen for each 100 square feet. If you soil has more than 5% organic matter you will only need to apply ½ ounce of nitrogen per 100 square feet at the beginning of the season.

<b>Table A: Nitrogen Sources</b>				
<b>Typical Nitrogen Fertilizers</b>	<b>Fertilizer Analysis</b>	<b>Amount of this fertilizer needed to supply 3.5 ounces of nitrogen per 100 square feet</b>	<b>Rate of nutrient release</b>	<b>Other considerations</b>
Urea (synthetic)	46-0-0	7.5 oz.	Rapid	Inexpensive, concentrated, may be difficult to apply
Calcium nitrate (inorganic)	15-0-0	23 oz.	Rapid	can help raise pH
Ammonium sulfate (inorganic)	21-0-0	16.5 oz	Medium	Can help lower pH, can use for acid loving crops
Blood Meal (natural organic)	12-0-0	29 oz.	Medium-rapid	Commonly available
Corn Gluten (natural organic)	10-0-0	35 oz.		Expensive
Cotton Seed Meal (natural organic)	6-2-1	58 oz.	Slow-medium	Can help lower pH, also contains phosphorus and potassium
Sodium Nitrate (natural organic)	16-0-0	22 oz.	Rapid	Avoid high rates

When possible, apply about 1/3 of the nitrogen pre-plant, another 1/3 at transplanting possibly in the form of a starter solution and the rest a few weeks after planting as a side dress.

**For established vegetable gardens,** if nitrogen was not applied before planting, apply half the recommended rate as a side dress immediately and then the other half after a couple weeks.

Apply the side dress fertilizer 2 to 4" from the growing plant to avoid contact with green plant tissue, lightly rake into the soil and then water it in to move fertilizer through soil to where roots are growing.

Crop	Timing of Side dressing	Amount of Fertilizer to Apply per 100 sq. ft. *			
		Urea	Calcium Nitrate	Ammonium Nitrate	Ammonium Sulfate
		46-0-0	16-0-0	33-0-0	20-0-0
Asparagus	End of harvest	¼ cup	½ cup	¼ cup	½ cup
Beets, Carrots, Celery, Endive, Turnips	When 4" tall	¼ cup	½ cup	¼ cup	½ cup
Cabbage, Cauliflower	Leaves 4-6" tall	¼ cup	½ cup	¼ cup	½ cup
Cucumbers, Melons, Pumpkins, Squash	When vines spread	¼ cup	½ cup	¼ cup	½ cup
Eggplant, Peppers, Tomatoes	Fruit set	¼ cup	½ cup	¼ cup	½ cup
Potatoes	When 8" tall	¼ cup	½ cup	¼ cup	½ cup
Spinach	When 3" tall	¼ cup	½ cup	¼ cup	½ cup
Sweet corn	When 8" tall	¼ cup	½ cup	¼ cup	½ cup

### Phosphorus (P)

Phosphorus is important in early root development and improves the quality of vegetables and fruits.

Phosphorus does not move in the soil much. It is best to incorporate P into the soil before planting so the roots can access more of this nutrient. A starter phosphorus solution is often applied when transplanting small plants in early spring when soils are cold even when P levels are adequate.

If only P is needed, use Table C to help determine the amount of fertilizer to apply. If the garden is already established, select a water-soluble fertilizer with a high percentage of P and apply according to label directions.

**Animal manures**, fresh and composted, often contain high levels of P. Regularly adding manure to gardens can raise soil P to excessive levels.

Soil Test Level and Recommended Amount (oz) of Phosphate to Apply /100 sq. ft.		Some Fertilizer Sources that contain Phosphorus			
		Inorganic fertilizers		Organic fertilizers	
Soil Test Level	Prior to Planting apply phosphate (P <sub>2</sub> O <sub>5</sub> )	Super-phosphate	Triple Super-phosphate	Bone Meal	Rock phosphate
		0-20-0	0-46-0	1-15-0 also contains N	0-3-0
		Quickly available	Quickly available	Slowly available	Slowly available
Amount of product to apply per 100 square feet					
Very low	6 oz.	30 oz.	13 oz.	2½ lbs.	12½ lbs.
Low	4.5 oz.	22 oz.	10 oz.	1.8 lbs.	9¼ lbs.
Medium	3 oz.	15 oz.	6.5 oz.	1¼ lbs.	6¼ lbs.
High	1.5 oz.	8 oz.	8 oz.	.6 lbs.	3 lbs.

## Potassium (K)

Potassium increases water efficiency, improves stress tolerance and the quality of seeds and fruit. Some of the potash (a source of potassium) should be applied before planting, mixed into the soil and the rest applied at the time of planting. If only K is needed use Table D to help determine the amount of fertilizer to apply.

Table D: Soil Test Levels and Pre-Plant Potassium recommendations					
Soil Test Level and Recommended Amount of Potash to Apply /100 sq. ft.		Fertilizer Sources that contain Potassium (K)			
		Inorganic fertilizers	Organic fertilizers		
Soil Test Level	Prior to Planting apply potash (K <sub>2</sub> O)	Potash 0-0-60	Kelp 1-0-2	Greensand	Wood Ashes*
		60% K <sub>2</sub> O	2% K <sub>2</sub> O	7% K <sub>2</sub> O	3-7% K <sub>2</sub> O
		Quickly available	Quickly available	Very slowly available	Quickly available
Amount of product to apply per 100 square feet					
Very low	4 oz.	6.5 oz.	12½ lbs.	3½ lbs.	8¼ - 3½ lbs.
Low	3 oz.	5 oz.	9.3 lbs.	2½ lbs.	6¼ - 2½ lbs.
Medium	2 oz.	3 oz.	6¼ lbs.	1¾ lbs.	4 - 1¾ lbs.
High	1 oz.	1.5 oz.	3 lbs.	.8 lb.	2 - ¾ lbs.

\* Wood ashes are not used as an amendment to improve soil physical properties. However, they do contain potassium and can be used to raise soil pH in place of lime. Do not apply more than 10 lbs. of wood ashes per 100 square feet and avoid applying both lime and wood ashes as these actions can raise the soil pH beyond the optimum range of 6.0-7.0. When using wood ashes as a lime substitute, be sure to incorporate them into the upper 4 to 6 inches of soil in advance of seeding or transplanting.

### Matching Your Soil Test Levels with a Fertilizer Ratio

Notice the proportion of nutrients recommended on the soil test report. **If all nutrients** are at a low or medium level, a fertilizer with a 1:1:1 ratio, for example: 10-10-10, 18-18-18, 20-20-20, could be ideal. The exact amount to apply would be different based on the concentration of the fertilizer.

When the exact fertilizer ratio desired is not readily available select a fertilizer close to the needed ratio of nutrients, remembering a little less may be better than too much. Table E provides some possible fertilizer ratios based on soil test levels.

<b>Table E: Matching Your Soil Test Levels with a Fertilizer Ratio</b>				
<b>Soil Test Level</b>		<b>Fertilizer Ratio</b>	<b>Examples N-P-K</b>	<b>Approximate amount of fertilizer needed for a 100 square foot area</b>
<b>Phosphorus (P)</b>	<b>Potassium (K)</b>			
Very low	Very low	1:2:2	5-10-10	4 lbs.
Very low	Medium	1:2:1	5-10-5	4 lbs.
Low	Low	1:2:2	5-10-10	4 lbs.
Low	Medium	1:2:1	8-16-8	2.5 lbs.
Medium	Low	1:1:2		2.5 lbs.
Medium	Medium	1:1:1	10-10-10	2 lbs.
High*	Low	1:0:2		Use Table E to find K source
High*	Medium	1:0:1	19-0-19	Use Table E to find K source
Low	High*	1:2:0		Use Table D to find P source
Medium	High*	1:1:0		Use Table D to find P source

\* A recommendations will only be made if nutrients are low or very low.

**Re-test your soil every 2-3 years!**

More gardening information at [gardening.cornell.edu](http://gardening.cornell.edu)

version 11.13.12